



A Critical Review of Factors and Challenges Influencing Non-Medical Prescribers in Primary and Urgent Treatment Care Facilities in England

Mithun Thampi¹ and Biju Mathew^{2*} 

¹Advanced Clinical Practitioner, Kingswood Medical Centre, Basildon-Essex, UK

²Faculty of Health Sciences, Anglia Ruskin University, Chelmsford, UK

*biju.mathew@aru.ac.uk (Corresponding Author)

ARTICLE INFORMATION

Keywords:

Non-medical prescribers, Minor illness, Urgent care, Primary care, Barriers and Facilitators

ABSTRACT

Background: Non-medical prescribing in England has empowered allied health professionals such as nurses, paramedics, and pharmacists working in acute and primary care settings to prescribe within their competency area in suitable work settings. The opportunities for non-medical staff to prescribe in specific clinical areas of expertise have increased substantially and continue to do so. Prescribing is now an integral part of advanced clinical practice, which is not limited to just nurses but has expanded to other allied health professionals.

Purpose: To analyse the factors and challenges influencing prescribing for non-medical prescribers, during consultations with patients in primary and urgent treatment centre facilities

Methods: This study employed a critical systematic review of relevant articles chosen from electronic databases including CINAHL, Medline, Cochrane library, and Scopus. Keywords were used to formulate a search pattern using Boolean operators and suitable qualitative studies relevant to the inclusion criteria were selected

Results: Thematic analysis of the articles concluded that there were a lot of sub-themes including a lack of protocols to support decision-making, peer support, and difficult patients which were interdependent posing as a potential barrier or acting as a facilitator in certain consultations

Conclusion: The findings have provided adequate reassurance that non-medical prescribers were aware of the facilitators and barriers to non-medical prescribing. Organizational support and continued professional development are key components of the barriers and enablers for non-medical prescribers working in urgent and primary care settings.



DOI: [10.15415/jmrh.2022.82004](https://doi.org/10.15415/jmrh.2022.82004)

1. Introduction

Non-medical prescribing (NMP) was adopted into the United Kingdom's national healthcare system in 1989 (Department of Health, 2006). A lot of reforms have happened since its inception over 2 decades ago, and the benefits of empowering allied health professionals with independent prescribing have explored new ways of working (Health Education England, 2015). Introducing prescribing into their practice has allowed allied health professionals to be more clinically challenged in their roles, further enabling autonomy, and enhancing job satisfaction. However, the medical fraternity interpreted this approach with caution and concern that they were more likely to make mistakes (Day, 2005). This enhanced clinical responsibility invoked the need for non-medical prescribers to thoroughly understand the consultation process to improve their clinical and diagnostic reasoning skills and enable independent decision-making

outcomes (Franklin, 2017). Prescribing decisions form one of the cornerstones of patient safety and is a complex and challenging aspect of advanced practice. Practitioners with prescribing abilities must use it in the best interest of patients using practically the same prescribing rights, held by medical counterparts (Pearce, 2016). The future of the health force in the United Kingdom (UK) relies on workforce transformation and recognizing contributions from these highly trained and skilled multi-disciplinary professionals. The 2019 NHS long-term plan has highlighted the benefits of non-medical prescribing in advanced clinical practice. Extending prescribing responsibilities to allied healthcare professionals have enabled them to deal with minor illnesses and injuries that might need pharmacological interventions, without having to refer them to a medical modality, releasing time for medical clinicians to engage in complex medical decisions which will not form the remit of advanced non-medical practitioners (NHS England, 2019).

NMP has developed to include other allied professionals not just in the UK but also in the United States of America, Canada, New Zealand, Finland, Sweden, and Republic of Ireland. In the majority of these countries, they work collaboratively with supervising medical clinicians to deal with patients presenting with pre-specified conditions and established diagnoses (Stewart et al., 2012) with the introduction of prescribing rights for healthcare professionals. In this article, we focus on the education, training and practice of nonmedical prescribers in the UK. There are around 20,000 nurse independent prescribers, 2400 pharmacist supplementary/independent prescribers, several hundred allied health professional supplementary prescribers and almost 100 optometrist supplementary/independent prescribers. Many are active prescribers, managing chronic conditions or acute episodes of infections and minor ailments. Key aims of nonmedical prescribing are as follows: to improve patient care; to increase patient choice in accessing medicines; and to make better use of the skills of health professionals. Education and training are provided by higher education institutions accredited by UK professional bodies/regulators, namely, the Nursing and Midwifery Council, General Pharmaceutical Council, Health Professions Council and General Optical Council. The programme comprises two main components: a university component equivalent to 26 days full-time education and a period of learning in practice of 12 days minimum under the supervision of a designated medical practitioner. Course content focuses on the following factors: consultation, decision making, assessment and review; psychology of prescribing; prescribing in team context; applied therapeutics; evidence-based practice and clinical governance; legal, policy, professional and ethical aspects; and prescribing in the public health context. Nonmedical prescribers must practise within their competence, demonstrating continuing professional development to maintain the quality engendered during training. Despite the substantial progress, there are several issues of strategy, capacity, sustainability and a research evidence base which require attention to fully integrate nonmedical prescribing within healthcare.

DOI: 10.1111/j.1365-2125.2012.04204.x
ISSN: 1365-2125
issue: 4
journalAbbreviation: Br J Clin Pharmacol
language: eng
note: PMID: 22300374
nPMCID: PMC3477334
page: 662-667
source: PubMed
title: Educating nonmedical prescribers
volume: 74
author: [{"family": "Stewart", "given": "Derek"}, {"family": "MacLure", "given": "Katie"}, {"family": "George", "given": "Johnson"}]
issued: {"date-parts": [[2012, 10]]}
schema: <https://github.com/citation-style-language/schema/raw/master/csl-citation.json>

As non-medical prescribers

are required to have many years of experience engaging with patients before beginning their prescribing training, it is reasonable to assume that this experience, together with specialised prescribing training and evaluation of competency, will result in a safer and more appropriate prescribing (McIntosh et al., 2016). Integrated care was the proposal put forward to bring in improved ways of working by challenging traditional pathways and using the most appropriate healthcare professional for the task at hand. The need for patient-centred care in a responsive approach in urgent care settings was highlighted long before from 2014 in the Five years NHS forward view (NHS England, 2014). Adequately trained NMPs were integral to these developments. Health Education England (Health Education England, 2015) proposed the benefits of non-medical prescribing in primary care to support medicine optimisation for long-term patients hence streamlining care for patients. NMPs who have undertaken enhanced qualifications in consultation and assessment are also responsible for diagnosis and prescribing for a wide range of acute, chronic, and complex conditions in urgent and primary care settings (Graham-Clarke et al., 2019) through extending healthcare professionals' roles. More recent government health service policy focuses on the increased demand and the need for efficiency. This systematic policy review aimed to describe any changes in government policy position and the role that non-medical prescribing plays in healthcare provision.

METHOD: The systematic policy review included policy and consultation documents that describe independent non-medical prescribing. A pre-defined protocol was registered with PROSPERO (CRD42015019786). The benefits of non-medical prescribing by advanced clinical practitioners have been highlighted in numerous studies and reports. However, the challenges and factors faced by NMPs in Urgent care settings are yet to be fully understood (Courtenay et al., 2017) employer, where the prescribing qualification is used, care setting and service provided. The specific question of prescribing behaviour and the challenges faced by NMP, especially in consultations with patients with minor illnesses is even less explored (Nelson et al., 2019) understand common implementation barriers, and identify measurable impacts or unintended consequences.

Design and setting In-depth qualitative comparison of three role initiatives in general practices in one area of Greater Manchester, England; that is, advanced practitioner and physician associate training schemes, and a locally commissioned practice pharmacist service.

Method Semi-structured interviews and focus groups with a purposive sample of stakeholders involved in the implementation of each role initiative were conducted. Template analysis enabled the production of pre-determined and researcher-

generated codes, categories, and themes.\nResults The final sample contained 38 stakeholders comprising training/service leads, role holders, and host practice staff. Three key themes captured participants' perspectives: purpose and place of new roles in general practice, involving unclear role definition and tension at professional boundaries; transition of new roles into general practice, involving risk management, closing training-practice gaps and managing expectations; and future of new roles in general practice, involving demonstrating impact and questions about sustainability.\nConclusion This in-depth, in-context comparative study highlights that introducing new roles to general practice is not a simple process. Recognition of factors affecting the assimilation of roles may help to better align them with the goals of general practice and harness the commitment of individual practices to enable role sustainability.”,”container-title”：“British Journal of General Practice”,”DOI”：“10.3399/bjgp19X704117”,”ISSN”：“0960-1643, 1478-5242”,”issue”：“684”,”journalAbbreviation”：“Br J Gen Pract”,”language”：“en”,”license”：“© British Journal of General Practice 2019. This article is Open Access: CC BY-NC 4.0 licence (<https://creativecommons.org/licenses/by-nc/4.0/>). The scope of this study will be limited to systematically reviewing articles exploring the influences on prescribing decisions by various health professionals including advanced nurses, pharmacists and paramedics working in secondary settings, urgent and primary care settings.

The focus of this review will be on non-medical prescribers, therefore articles pertaining to this specific topic have been considered for review. The studies chosen for this have originated from the United Kingdom as they offer in-depth analysis and the impact across all areas where non-medical prescribers independently review patients presenting with minor illnesses. The various gaps in knowledge and how it affects current practice will also be explored. The literature search will be discussed, and the chosen articles will be critically reviewed explaining the methodology and also the limitations. Recommendations for future research will also be highlighted and discussed.

1.1. Background

This chapter gives a descriptive review of the larger context of the obstacles and enablers affecting non-medical prescribers in primary and urgent care settings. The concept of primary care networks, non-medical prescribing and independent prescribing will be discussed to provide a wider perspective. There is a large body of literature on the prescription behaviour of non-medical prescribers, but very few studies examine this topic in the context of primary and urgent care consultations.

1.2. Primary Care Networks

The prevalence of chronic diseases like diabetes and heart disease, as well as mental health problems, means that more individuals may need to make use of their community healthcare facilities. To satisfy these demands, practices have started forming primary care networks with their local communities, mental health services, social care providers, pharmacies, hospitals, and volunteer agencies (The King's Fund, 2019). One of the most consequential practical attempts to maintain the NHS's survival in a time of major change and a desire for improved efficiency is the expanding role of the non-medical prescriber (NHS England, 2018). Independent prescription (IP) by nurses, pharmacists, and allied health professionals is a crucial element of workforce transformation in the UK healthcare system, particularly in primary care networks, in order to address staff shortages and the increased demand for pharmaceutical interventions (Edwards et al., 2022). Since its adoption, the role of prescribing for nurses and pharmacists has constantly expanded and evolved, significantly improving patient care and cost-effectiveness (Magowan, 2020). The aim of this innovation was to free up medical practitioners consulting time to focus on more complex cases, thus reducing waiting times for appointments and preventing further harm to patients presenting with minor illnesses (NHS England, 2018). However, Hacking and Taylor (2010) have mentioned that clinicians who were assisting NMPs with their clinical decisions and prescribing outcomes had a sizeable time and commitment added to their work.

1.3. Non-medical Prescribing and Independent Prescribing

The conventional doctor-led prescribing format did not keep up with the growing demands in primary care, hence seeking out new methods of prescribing by allied professionals, became necessary to sustain patient access to prescription medications (Armstrong et al., 2021)written in English language. Exclusion criteria: research conducted in secondary care or outside the UK.\n\nFindings:\n\n285 studies were identified; 15 were eligible for critical appraisal. Key themes were: NMP's positive perceptions were autonomy, job satisfaction and colleague support; negative perceptions included risk, lack of continuous professional development (CPD). The number of NMPs in England, with independent prescribing rights, has significantly risen in the last decade. Courtenay et al approximately estimate that over 90,000 independent non-medical prescribers are practising in the United Kingdom, and exercising their prescribing rights to clinically appropriate presentations(Courtenay et al., 2017) employer, where the prescribing qualification is used, care setting and service provided. However, Wider adoption of

nonmedical prescribing practice is frequently hindered by local regulatory obstacles and opposition from the medical community, which has raised concerns about professional autonomy, patient safety, the diagnostic competency of nonmedical prescribers, and costs (Robertson, 2022). There are two kinds of NMP in the United Kingdom: supplemental prescribing (SP) and independent prescribing (IP). SP, which was launched in 2003, is described as a “voluntary collaboration between an independent prescriber and a supplemental prescriber to execute an agreed patient-specific clinical management plan with patient consent” (Stewart et al., 2017) consideration of two key challenges. The review identified seven systematic reviews of influences on prescribing decision-making, processes of prescribing, and barriers and facilitators to implementation. Decision making was reported as complex with many, and often conflicting, influences. Facilitators of NMP included perceived improved patient care and professional autonomy, while barriers included lack of defined roles and resource pressures. Three systematic reviews explored patient outcomes that were noted to be equivalent or better to physician prescribing. In particular, a Cochrane review of 46 studies of clinical, patient-reported, and resource-use outcomes of NMP compared with medical prescribing showed positive intervention-group effects. Despite positive findings, authors highlighted high bias, poor definition and description of ‘prescribing’ and the ‘prescribing process’ and difficulty in separating NMP effects from the contributions of other healthcare team members. While evidence of benefit and safety is essential to inform practice, for NMP to be implemented and sustained on a large scale, there needs to be clear commitment at the highest level. The approach being taken by the Scottish Government to pharmacist prescribing implementation may inform developments in other professions and countries. The vision is that by 2023, all pharmacists providing pharmaceutical care will be pharmacist-independent prescribers. There are, however, challenges to implementing NMP into working practice; two key challenges are the need for sustainable models of care and evaluation research. These challenges could be met by considering the theoretical basis for implementation, and robust and rigorous evaluation.”, “container-title”: ”Therapeutic Advances in Drug Safety”, ”DOI”: ”10.1177/2042098617693546”, ”IS SN”: ”2042-0986”, ”issue”: ”6”, ”journalAbbreviation”: ”Ther Adv Drug Saf”, ”note”: ”PMID: 28607668\nPMCID: PMC5455843”, ”page”: ”183-197”, ”source”: ”PubMed Central”, ”title”: ”Future perspectives on nonmedical prescribing”, ”volume”: ”8”, ”author”: [{“family”: ”Stewart”, ”given”: ”Derek”}, {“family”: ”Jebara”, ”given”: ”Tessime”}, {“family”: ”Cunningham”, ”given”: ”Scott”}, {“family”: ”Awaisu”, ”given”: ”Ahmed”}, {“family”: ”Pallivalapila”, ”given”: ”Abdulr”}], ”issued”: {“date-parts”: [[“2017”, 6]]}], ”schema”: ”https://github.com/citation-style-language/schema/raw/master/csl-citation.json”} . SP designation is now conferred to those who qualify as independent prescribers. The introduction of IP in 2006 was a later development (Courtenay & Griffiths, 2022) and described as “prescribing” by a practitioner (e.g., doctor, dentist, nurse, pharmacist) responsible and accountable for the evaluation of patients with undiagnosed or diagnosed diseases and for choices about clinical care, including prescribing.

It has been determined that allied health professionals such as physiotherapists, podiatrists, nurses, paramedics, pharmacists, and radiographers play a crucial role in this essential transformative shift (Edwards et al., 2022). NMPs are required to address these gaps as a result of the decline of Practitioners (GP) and the changing methods of working in primary care (Winter, 2019). Independent prescribing by paramedics working in advanced roles has been adopted into the prescribing remit for NMPs since 2018, with the exclusion of prescribing controlled drugs (NHS England, 2018).

1.4. Statement of the Problem

The influence of NMP on patients, practitioners and organisations has been the focus of many studies on the topic so far. Job satisfaction, increased autonomy, and opportunities to improve patient care have been the main factors associated with a rise in prescribing practice (Courtenay et al, 2018). The competency framework devised by the Royal Pharmaceutical Society (Royal Pharmaceutical Society, 2016) specifies the requirements of all prescribers and supersedes the previous guidelines by National Institute for Healthcare Excellence (NICE, 2012) enabling all allied practitioners, who have satisfactorily completed an approved prescribing course to prescribe within their scope of practice. Despite this solid framework, there are still substantial variations in how independent prescribers use their prescribing qualifications, with some preferring only to prescribe in certain circumstances or not at all (Nuttall, 2018). Due to an ageing population, complicated polypharmacy requirements, and an increase in co-morbidities, all prescribers face significant challenges. To stay current on new medications and potential interactions, non-medical prescribers must actively develop and maintain their competence to prescribe (Royal Pharmaceutical Society, 2016), which can extend beyond the confinements of their domain of expertise. However, the challenges and factors influencing non-medical prescribers are yet to be fully understood. Prescribing within limits of practice and knowledge is appropriate to newly qualified prescribers who

are experienced practitioners, but new to the prescribing remit (Stewart et al., 2017) consideration of two key challenges. The review identified seven systematic reviews of influences on prescribing decision-making, processes of prescribing, and barriers and facilitators to implementation. Decision making was reported as complex with many, and often conflicting, influences. Facilitators of NMP included perceived improved patient care and professional autonomy, while barriers included lack of defined roles and resource pressures. Three systematic reviews explored patient outcomes that were noted to be equivalent or better to physician prescribing. In particular, a Cochrane review of 46 studies of clinical, patient-reported, and resource-use outcomes of NMP compared with medical prescribing showed positive intervention-group effects. Despite positive findings, authors highlighted high bias, poor definition and description of 'prescribing' and the 'prescribing process' and difficulty in separating NMP effects from the contributions of other healthcare team members. While evidence of benefit and safety is essential to inform practice, for NMP to be implemented and sustained on a large scale, there needs to be clear commitment at the highest level. The approach being taken by the Scottish Government to pharmacist prescribing implementation may inform developments in other professions and countries. The vision is that by 2023, all pharmacists providing pharmaceutical care will be pharmacist-independent prescribers. There are, however, challenges to implementing NMP into working practice; two key challenges are the need for sustainable models of care and evaluation research. These challenges could be met by considering the theoretical basis for implementation, and robust and rigorous evaluation." "container-title": "Therapeutic Advances in Drug Safety", "DOI": "10.1177/2042098617693546", "ISSN": "2042-0986", "issue": "6", "journalAbbreviation": "Ther Adv Drug Saf", "note": "PMID: 28607668\nPMCID: PMC5455843", "page": "183-197", "source": "PubMed Central", "title": "Future perspectives on nonmedical prescribing", "volume": "8", "author": [{"family": "Stewart", "given": "Derek"}, {"family": "Jebara", "given": "Tessime"}, {"family": "Cunningham", "given": "Scott"}, {"family": "Awaisu", "given": "Ahmed"}, {"family": "Pallivalapila", "given": "Abdulrouf"}, {"family": "MacLure", "given": "Katie"}], "issued": {"date-parts": [[2017, 6]]}, "schema": "https://github.com/citation-style-language/schema/raw/master/csl-citation.json"} .

Anti-microbial resistance (AMR) is a global public health threat and overuse is the main driver for the development of drug-resistant pathogens (WHO, 2021). The consequence of increased drug resistance is that infections are becoming increasingly difficult or often impossible to treat. 80% of antimicrobial prescribing occurs in primary care settings, where non-medical prescribers

from nursing and other allied health sectors are actively involved in prescribing decisions (Beech, 2015). A growing number of non-medical prescribers often inappropriately prescribing antibiotics can potentially cause an increase in this resistance. Responsible and appropriate prescribing of anti-biotics will require commitment and engagement from the wider group of prescribers including non-medical prescribers (Courtenay et al., 2017) employer, where the prescribing qualification is used, care setting and service provided. As the number of prescribing nurses, pharmacists, and AHCPs grows at an ever-increasing rate, problems regarding access to Continuing Professional Development support, and organisational infrastructure continue to slow down the acceptance of non-medical prescribing practice (Nuttall, 2018). Hence further research is required to enable NMPs to grow and extend their scope and remit of prescribing practice, and for this crucial service to thrive. The bulk of support for prescribing activity comes from colleagues and other members of the multidisciplinary team, as well as resources such as NICE Clinical Knowledge Summaries and a variety of programme-integrated support tools (Hubbard, 2019). However, there are challenges with evidence-based practice, as well as independent factors that influence its use in Non-Medical Prescribing. Evidence that is of poor quality, fragmentary, or inconsistent has an impact on the quality of the scientific data and hence may affect NMP (Booth, 2016). The technique of utilising evidence by NMPs must be more fluid and dynamic (Griffith & Tengnah, 2014) patients, their employers and the profession, so it is vital you have a clear understanding of the legal, ethical and professional dilemmas you will face in the course of your career. This book introduces the legal and professional requirements of safe nursing in clear, straightforward terms and helps you to understand how they apply to nursing practice. The third edition of this popular book has been fully updated with changes to the law and professional requirements, and includes new case studies, scenarios and activities from all fields of practice and a clearer colour text design. Key Features:

- Each chapter is linked to relevant NMC Standards and Essential Skills Clusters so you can see what is required in order to become a registered nurse
- Scenarios and case studies show how the law applies to your nursing practice
- Activities help you to build core skills such as critical thinking and reflection." "ISBN": "978-1-4462-7640-2", "language": "en", "note": "Google-Books-ID: KqyNAgAAQBAJ", "number-of-pages": "289", "publisher": "Learning Matters", "source": "Google Books", "title": "Law and Professional Issues in Nursing", "author": [{"family": "Griffith", "given": "Richard"}],

"family": "Tengnah", "given": "Cassam"}], "issued": {"date-parts": [[["2014", 1, 9]]]}}, "schema": "https://github.com/citation-style-language/schema/raw/master/csl-citation.json"} . Another criticism of evidence-based practice is its frequently irrelevant connection to clinical governance (Lim et al., 2017).

Smith et al. (2014) found that the challenges and enablers to prescribing for NMPs operating in Primary care and the acute sector were comparable. This is counterintuitive as NMPs working in primary care are nearly often direct employees of the General Practice and the company they operate, but NMPs working in secondary care are part of major NHS organisations working with standardised protocols and agreed frameworks (Bryant-Lukosius et al., 2016) stakeholders identified the need for guidance to generate strategic evaluation data. This article describes an evaluation framework developed to inform decisions about the effective utilization of APN roles across the country.

nAPPROACH: A participatory approach was used by an international group of stakeholders. Published literature and an evidenced-based framework for introducing APN roles were analyzed and applied to define the purpose, target audiences, and essential elements of the evaluation framework. Through subsequent meetings and review by an expert panel, the framework was developed and refined.

nFINDINGS: A framework to evaluate different types of APN roles as they evolve to meet dynamic population health, practice setting, and health system needs was created. It includes a matrix of key concepts to guide evaluations across three stages of APN role development: introduction, implementation, and long-term sustainability. For each stage, evaluation objectives and questions examining APN role structures, processes, and outcomes from different perspectives (e.g., patients, providers, managers, policy-makers. In addition to helping administrators strategically plan for the growth of NMPs and the improvement of the infrastructure to accommodate this growth, gaining a deeper understanding of the unique challenges encountered by primary care NMPs may open up new doors of opportunity for practitioners in this field (Park et al., 2018). According to Harvey & Kitson (2015), holistic patient care may be achieved by integrating several factors, including evidence, experience, the viewpoint of patients, and the use of available resources (Harvey & Kitson, 2015). However, the emphasis of training for nurses and other non-medical prescribers should be on enhancing their confidence and ability to treat patients in the urgent care setting without relying solely on pharmaceutical interventions (Rowbotham et al., 2012) general practitioners frequently prescribe antibiotics, citing non-clinical reasons such as patient pressure. Nurses increasingly manage people with respiratory tract infections,

but research has not yet explored their experiences within such consultations.

DESIGN: Semi-structured interviews and focus groups.

METHODS: Fifteen semi-structured interviews and three focus groups ($n=5$, $n=4$, and $n=12$). The main scope of this study is independent prescribing behaviour by non-medical prescribers, therefore articles pertaining to prescribing by independent prescribing NMPs in primary and urgent care settings will be considered for review. This review will attempt to identify the themes influencing prescribing practice in urgent and primary care settings during minor illness presentations. This study was conducted in accordance with strict ethical ideals and principles. As stated by Beauchamp & Childress (2019), the researcher adhered to stringent ethical principles and ideals while conducting this investigation.

2. Methodology

This section includes a summary of several techniques, research designs, and paradigms. It emphasises the essential strategies utilised in establishing the assessment of the variables and obstacles affecting prescribing by non-medical prescribers during consultations with patients with a variety of minor illness presentations in primary and urgent care facilities. This section will describe the methodology along with the strategy used to collect, examine, and evaluate the methodology of the articles included in the research. The purpose of this section is to build a rigorous procedure for verifying and producing reliable data, which will be addressed in the succeeding sections. This established the groundwork for later sections that addressed the research question.

2.1. Research Question

A critical literature review must begin with a query, as well as a defined purpose and reason (Polit & Beck, 2017). Developing a clear and succinct purpose is essential when determining a research topic (Booth & Carroll, 2015) and it is the best approach for any literature review. The specific research question of this review is: What are the factors and challenges influencing prescribing by non-medical prescribers during consultations involving patients with various minor illness presentations in primary and urgent treatment care facilities.

2.2. Aims and Objectives

The study's goals and objectives should be stated in a way that the reader may grasp them quickly and readily (Kumar, 2010), while also making sure that they are clear and related to the research issue (Offredy & Vickers, 2010). The aim

of this literature review is to identify specific challenges faced by NMPs, particularly in primary and urgent care facilities. This information may be utilised to empower the prescribing behaviour of NMPs by strengthening and supporting the practice of independent prescribing in order to promote patient-centred, safe, effective, and efficient pharmacological treatments during patient consultations.

2.3. Specific Objectives

1. Formulate a search strategy to identify pertinent research articles on the factors influencing NMP prescribing behaviour in urgent and primary care consultations.
2. Conduct a critical review that will identify specific challenges faced by NMPs during patient consultations.
3. Discuss the implications of the themes and sub-themes identified during the critical review of the chosen articles.
4. Explore the current strategies and interventions in place to support NMPs to effectively deal with challenges in prescribing practice through systematic review of the chosen articles.
5. To discuss the implications of the findings to support the suggestions to enable NMPs to confront the determinants and barriers impacting prescribing behaviour.

2.4. Search Strategy

The entirety of a systematic review's quality is decided by the breadth and calibre of the literature found during the search (Gerrish & Lacey, 2013). A high-quality, systematic, critical literature review incorporates broad search tactics to guarantee the inclusion of the largest proportion of the most relevant original research publications available (Aveyard, 2018). Clear documentation of the search strategy ensures that the process is methodical, unambiguous, and repeatable (Greenhalgh, 2010). The Population, Exposure, and Outcome (PEO) framework was used as shown in table 1 below to develop keywords and to search for qualitative research publications pertaining to the determinants and obstacles impacting prescription behaviour in NMPs in Urgent care consultations (Bettany-Saltikov & McSherry, 2016).

Table 1: PEO framework.

PEO Criteria	Keywords
Population	Non-medical prescribers in prescribing settings
Exposure	Minor illness, Urgent care, and primary care
Outcome	Barriers and Facilitators

This tool is a basic technique for formulating the search query despite critiques over its inability to locate relevant literature materials (Brun, 2013). Aveyard et al., (2021) highly readable guide to how to undertake a literature review in health and social care, tailored specifically for postgraduate study. Essential reading for all those undertaking any study at post-graduate level, the book provides clarity and a step by step approach to doing a literature review from start to finish which will enable you to:

- Identify which type of review is appropriate for your study
- Select the literature that you need to include in your review
- Search for, appraise and analyse relevant literature
- Write up your reviewCrucially the book explores the common features of a broad range of types of literature review, which serve different functions – including the literature review that is a pre-requisite prior to a larger empirical study, and the literature review that is a study in its own right. With real-life examples of written research and succinct summaries at the end of each chapter, *A Post-Graduate's Guide to Doing a Literature Review in Health and Social Care* is the ideal text for students wanting to get the very most from their study.”, “ISBN”：“978-0-335-24992-3”, “language”：“en”, “note”：“Google-Books-ID: AMkvEAAAQBAJ”, “number-of-pages”：“202”, “publisher”：“McGraw-Hill Education (UK) emphasised that one instrument is not preferable to another and that a single tool often cannot provide all the required literature. The search for qualitative publications is also challenging, labour-intensive, time-consuming, and difficult to repeat Booth (2016). The search was accomplished by combining topics and free terms using the “AND” and “OR” Boolean operators as shown in table-2 below. The fundamental words and their synonyms were included throughout all electronic databases, including CINHAL, SCOPUS, and Medline, to provide more targeted and prolific outcomes. During the search, truncations such as asterisks (*) and parentheses () were also used. As shown in Table 3 and 4 below, the search was limited to studies within the United Kingdom as that was the specific demographic being explored in the research question. Due to a lack of accessible resources to translate and understand them, the search was restricted to English-language articles only. There was a methodical search of databases for pertinent literature.

Table 2: Summary of the search phrases using Boolean operator.

	Population (P)	Exposure (E)	Outcome (O)
Boolean operator		and	and
	Non-medical prescribing	Facilitators	Urgent care
and	Non-medical prescriber	Barriers	Primary care
or	Non-medical independent prescriber	Challenges	Minor illness

2.5. Inclusion and Exclusion Criteria

Due to the paucity of published studies on the issue, primary research utilising qualitative studies addressing the facilitators


and obstacles to non-medical prescription was incorporated to broaden the scope and emphasis of the study.

Table 3: Inclusion Criteria.

1. Qualitative research
2. Dated within 10 years
3. Primary studies
4. English papers
5. Semi-structured Interviews

Table 4: Exclusion Criteria.

1. Quantitative research
2. Dated above 10 years.
3. Secondary studies.
4. Non-English papers.
5. Service evaluations.

**Figure 1:** PRISMA flow Diagram.

The huge volume of results from the database and search engine posed one of the greatest obstacles throughout the search for relevant material. To be able to assess the most relevant research, the author spent a great deal of time acquiring a broad range of subject-matter expertise.

Accessing journals and articles that required a subscription or payment was another obstacle. Access to many recent articles and publications was restricted, which posed a challenge to the ability to comprehend the most current information on the topic at hand. Following an exhaustive search of all databases, 148 articles were discovered. To ensure that the search was exhaustive and that no research was overlooked or excluded mistakenly, 13 more publications were uncovered using the reference lists of included papers and a general search (Parahoo, 2014) processes and issues. The author guides readers who are new to research but also introduces new debates and perspectives to those with some experience wanting to develop their skills further. This popular book equips students with the information and skills they need to read, comprehend and critique research. Whether an undergraduate taking an introductory research module, a postgraduate nursing student embarking on a project, or an experienced

practitioner wanting to sharpen your skills, Parahoo's accessible writing style will ensure readers are able to utilise research throughout their study and in everyday practice. New to this Edition:- Three new chapters on qualitative methods, introducing grounded theory, phenomenology and ethnography- Updated narrative and research examples to ensure content and application is relevant", "ISBN": "978-1-137-28127-2", "language": "en", "note": "Google-Books-ID: 0yNIEAAQBAJ", "number-of-pages": "437", "publisher": "Bloomsbury Publishing", "source": "Google Books", "title": "Nursing Research: Principles, Process and Issues", "title-short": "Nursing Research", "author": [{"family": "Parahoo", "given": "Kadet"}], "issued": {"date-parts": [[2014, 5, 30]]]}, "schema": "https://github.com/citation-style-language/schema/raw/master/csl-citation.json"} . 91 papers did not match the requirements for inclusion. After a careful critical evaluation of the selected publications, ten main studies pertinent to this literature review were selected. The PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) standards for Systematic Reviews (Moher et al., 2009) flowchart in Figure-1 depicts the comprehensive selection process of research identification, review, inclusion and exclusion.

Table-5: Characteristics and Themes from selected studies.

SL	Author and Year	Title and DOI	Design	Participant group	Themes	
					Facilitators	Barriers
1.	Louise Cope, Mary Tully, Jason Hall. 2019	An exploration of the perceptions of non-medical prescribers, regarding their self-efficacy when prescribing, and their willingness to take responsibility for prescribing decisions. https://doi.org/10.1016/j.sapharm.2019.05.013	Cross-sectional survey. Framework Analysis.	<i>Nurse, Pharmacist and physiotherapist NMP.</i>	Length of time qualified as NMP. The professional role of the NMP	Lack of confidence. Lack of teaching around clinical versus legal responsibility when prescribing
2.	Roisin Lennon and Anne Fallon. 2018	The experiences of being a registered nurse prescriber within an acute setting. https://doi:10.1111/jocn.14087 .	Semi-structured interviews.	<i>Nurse Independent prescribers.</i>	Enhanced and complete episodes of patient care. Improved service and safe prescribing. Autonomy and Increased job satisfaction.	Increased workload. Unawareness of role of the non-medical prescriber. Lack of recognition. Lack of resources.
3.	Samantha Rowbotham, Rosemary Lim, Sarah Peters, Kathryn Yates, Angel charter. 2019.	Examining influences on antibiotic prescribing by nurse and pharmacist prescribers: a qualitative study using the Theoretical Domains Framework and COM-B http://dx.doi.org/10.1136/bmjopen-2019-029177	Semi-structured interviews using the Theoretical Domains Framework and Capability, Opportunity and Motivation for Behaviour.	<i>Pharmacist and Advanced Nurse Practitioner Non-medical prescribers.</i>	Knowledge of current prescribing guidelines. Clinical examination skills.	Lack of confidence in newly qualified prescribers. Time limitations on consultations. Emotions (Tiredness, stress, empathy). Fear of complaints as a consequence of non-prescribing.

4.	Emma Graham-Clarke, Alison Rushton, John Marriott. 2021.	A Delphi study to explore and gain consensus regarding the most important barriers and facilitators affecting physiotherapist and pharmacist non-medical prescribing. https://doi.org/10.1371/journal.pone.0246273	Delphi technique using Questionnaires.	<i>Pharmacist and Physiotherapist Independent non-medical prescribers.</i>	Independent mode of working. Streamlined care for patients. Improved patient care-reduced waiting times.	Lack of support and time to develop skills. Lack of acquisition of clinical skills. Personal confidence in prescribing skills. Prescribing legislation and indemnity requirements.
5.	Nicky Wilson, Catherine Pope, Lisa Roberts, Robert crouch. 2020.	Limited pharmaceuticalisation: a qualitative case study of physiotherapist prescribing practices in an NHS Trust in England following the expansion of non-medical prescribing in the UK https://doi.org/10.1111/1467-9566.13050	Semi-structured Interviews.	<i>Physiotherapist Independent Non-medical prescribers.</i>	Delivering integrated care.	Lack of knowledge base. Lack of jurisdictional boundaries in prescribing for physiotherapist non-medical prescribers. Reduced Follow-up capacity. Time pressures. Lack of communication between the medical fraternity and non-medical prescribers.
6.	Robert Weglicki, Julie Reynolds, Peter Rivers. 2015.	Continuing professional development needs of nursing and allied health professionals with responsibility for prescribing. http://dx.doi.org/10.1016/j.nedt.2014.08.009	Semi-structured interviews.	<i>Nurse independent prescribers, Physiotherapists and Pharmacist non-medical prescribers.</i>	Optimise skills of the workforce within primary, acute and community sectors.	Personal Anxiety undermining confidence to prescribe. External factors including a lack of CPD training sessions. Need for support from medical and non-medical colleagues and clinical supervisors. Increased accountability and responsibility that prescribing confers upon NMPs.
7.	Maddox, C., Halsall, D., Hall, J. and Tully, M.P 2016.	Factors influencing nurse and pharmacist willingness to take or not take responsibility for non-medical prescribing https://doi.org/10.1016/j.sapharm.2015.04.001	Nurse and Pharmacist NMPs.	<i>Semi-structured Interviews using the critical incident technique.</i>	Improving access to medicines while maintaining patient safety.	Underpinning cautiousness. Risk of making errors. Vulnerable to exposure and criticism from regulators and the public. Lack of supportive clinical supervision. Lack of mentoring support from medical colleagues.
8.	Ali M. K. Hindi, Elizabeth M. Seston, Dianne Bell, Douglas Steinke, Sarah Willis. 2019	Independent prescribing in primary care: A survey of patients', prescribers' and colleagues' perceptions and experiences https://doi.org/10.1111/hsc.12746	Semi-structured interviews.	<i>Nurse Independent prescribers, Podiatrist and pharmacist non-medical prescribers.</i>	Competence and confidence to prescribe. Support from multi-disciplinary healthcare team. Cohesive team working. The convenience of independent prescribing for patients. Valuable addition to the team.	Lack of competence in certain areas. Inadequate Training. Organisational barriers due to workload pressures. Lack of independent awareness.

9.	Aseel Abuzour, Penny Lewis, Mary Tully. 2017	A qualitative study exploring how pharmacist and nurse-independent prescribers make clinical decisions https://doi.org/10.1111/jan.13375	Think-aloud methodology and semi-structured interviews.	<i>Nurse and Pharmacist independent prescribers.</i>	Opportunity for conjoint decision-making with patients. Better concordance to treatment along with health promotion and support. Contextual influences on clinical reasoning enabling individualised decision-making skills.	Lack of adequate knowledge in clinical assessment skills. Depending on other MDT to reach a treatment plan for prescribers not competent in certain assessment skills.
10.	Sally Jarman. 2022.	Describing prescribing identities: a qualitative study exploring non-medical prescriber identity https://doi.org/10.12968/jpp.2022.4.7.300	Semi-structured interviews.	<i>Non-medical prescribers including nurses, pharmacists and physiotherapists.</i>	Using role modelling to enhance own prescribing practice. Consolidating the new identity and adopting best practices from other NMPs. Greater sense of belonging.	Insecurities about the prescribing role about complaints from other modalities. Concerns that other health professionals will question their right to prescribe. Collectivity versus isolation in working. Increased level of anxiety about the new identity and perception by other health professionals.

3. Results

According to Kokosi & Harron, (2022) Synthesis is a type of analysis that contrasts and compares, requiring the author to work with two or more summaries found in the literature to produce a conclusion. Furthermore, they state that the author would employ data from multiple analyses to reach a different conclusion. Additionally, merging, synthesising, and extracting data from several research is a systematic review that goes beyond a standard literature review (Munn et al., 2014). Data analysis seeks to make sense of all gathered data and streamline the complex narrative (Williamson & Whittaker, 2017). Following data extraction, a narrative review utilising thematic analysis was conducted to identify recurring themes. The author of this systematic review intends to perform a data synthesis utilising theme analysis as part of this evaluation. According to Mays et al. (2005), thematic analysis is the most effective method for extracting themes from narrative reviews and bringing them together. The results of several qualitative research were combined to form a synthesis, and the themes emerged from the sub-themes. Thematic analysis is a way of doing qualitative research that is both thorough and comprehensive, and using this strategy helped the themes become more integrated and less fragmented (Parahoo, 2014). The chosen papers

were evaluated to see whether it had a clear description of the research's purpose, its relevance and significance, and whether the methodology and research design were adequate and justified. Clear and deliberate checks were performed to evaluate how themes were created from the data, as well as to determine whether the analysis was thorough enough, with adequate data to support the conclusions. The articles were also investigated to determine whether they included any contradicting data or whether there was a possibility of bias.

Table 6: Facilitators and Barriers Identified from Studies.

Facilitators	Barriers (Current limitations of non-medical person for clinical practice in England)
Conjoint decision-making.	Lack of confidence and adequate knowledge in consultation skills.
Autonomous mode of working.	Increased workload.
Improved service and safe prescribing.	Role clarity.
Use of role models/mentors.	Insecurity about the prescribing role.
Opportunities for advanced practice.	Reduced opportunities for follow-up.

3.1. Themes Identified from the Selected Studies

Theme 1: Confidence: This analytical theme demonstrates how crucial confidence in prescribing during the post-qualification transition phase for new prescribers. It also highlighted how critical it is for new prescribers who are already established clinicians in their field to have supervision and formal and informal support during the didactic phase. Cope et al. (2019) argue that the length of time qualified as an NMP has a positive influence on the confidence in prescribing; however, Lennon et al. (2018) argue that this can also pose a barrier by placing an increased workload on experienced prescriber practitioners. Rowbotham et al (Rowbotham et al., 2012) general practitioners frequently prescribe antibiotics, citing non-clinical reasons such as patient pressure. Nurses increasingly manage people with respiratory tract infections, but research has not yet explored their experiences within such consultations.\nDESIGN: Semi-structured interviews and focus groups.\nMETHODS: Fifteen semi-structured interviews and three focus groups (n=5, n=4, and n=12 illustrate “lack of confidence” in newly qualified prescribers as a major barrier, whereas Maddox et al. (2016) argue that personal emotions and fear of complaints as a result of non-prescribing are major barriers to prescribing in primary care settings. Most of the participants who participated in the research by Jarman, (2022) used different role models to assist them recognise their identities as prescribers. This role modelling was a major factor in boosting confidence especially in new prescribers. However, there were not enough good medical role models for some specialties and professions, especially when NMPs worked in a relatively new discipline of practice (Weeks et al., 2016).

Theme 2: External barriers: Demonstrating external barrier interdependence, the absence of coaching ties with physicians hindered the chance for informal assistance, prohibited prescription, and limited competency with certain medications or clinical situations (Graham-Clarke et al, 2021). In turn, this prompted the re-engagement of GP referral for prescribing and resulted in disproportionate patient medication management (Jarman, 2022).

Wilson et al. (2020) and Weglicki et al. (2015) discovered similar themes in their research about how external factors such as inadequate communication and a lack of follow-up arrangements put NMPs under pressure to use effective pharmacological interventions. Hindi et al. (2019), on the other hand, valued the reassurance of having approachable and available GPs for advice. However, according to Abuzour et al. (2017), the administrative staff in primary care practices played an important role in facilitating independent prescribing by directing appropriate patients to NMPs.

Theme 3: Competency: Access to formal, educational practise updates and resources, as well as the availability

of peer support, was acknowledged as vital for continuing prescribing proficiency (McHugh et al, 2020). According to Jarman, (2022), independent prescribers believed that NMPs could not successfully prescribe without extensive training and experience. This lack of training opportunities was especially felt by those participants who were unable to discuss prescribing dilemmas (Maddox, et al, 2016). As a result, they lost confidence in their abilities and were occasionally hesitant to prescribe. Hindi et al. (2019) note that NMPS were especially concerned about consultations that required them to prescribe outside of their clinical areas of expertise. Graham-Clarke et al. (2019) through extending healthcare professionals’ roles. More recent government health service policy focuses on the increased demand and the need for efficiency. This systematic policy review aimed to describe any changes in government policy position and the role that non-medical prescribing plays in healthcare provision.\nMETHOD: The systematic policy review included policy and consultation documents that describe independent non-medical prescribing. A pre-defined protocol was registered with PROSPERO (CRD42015019786, on the other hand, discussed how prescriptive authority improved daily clinical practice by improving expert and lateral thinking abilities and knowledge. These skills were not only useful to the NMPs, but they also meant that the patients received advanced care.

Theme 4: Improved Quality of care: This is the most universally occurring theme in most of the existing research on this topic. The provision of individualised, holistic treatment was regarded as fundamental to their prescribing responsibility (Berry et al, 2007). McIntosh et al. (2016) accentuates that prescribing was not considered as a task-based activity, but rather as a crucial component of their role that enabled NMPs to provide high-quality, patient-centred care and maximise health outcomes in the shortest time available. Adoption of non-medical prescribing is likely to result in a variety of improvements in patient care and professional practise (Rowbotham et al, 2019). Wilson et al. (2020) also highlight that non-medical prescribing enabled better continuity of care for patients discharged into the community with primary care follow-up, allowing NMPs to modify or tweak pharmacological interventions, saving time for both patients and secondary care providers.

Theme 5: Higher autonomy and Job satisfaction: Prescribing, according to the NMPs, enabled them to be independent, responsible, and accountable practitioners who delivered a comprehensive package of responsive and safe care to their service users (Hubbard, 2020). The sense of satisfaction, according to Lennon et al (2018), was evident in the feelings of pride and accomplishment expressed by the participants in their comments about what prescribing meant to them.

The increase in professional autonomy that resulted from prescribing authority was discussed favourably by nine of the participants in the research article on pharmacist and nurse-independent practitioner clinical prescribing decisions (Abuzour et al, 2017). According to Park et al. (2018), if a practitioner is truly accountable for their prescribing practise, they ought to be granted the required autonomy and freedom to make their own judgements by utilising the full remit of independent prescribing and should not be constrained in this endeavour by local regulatory guidelines. However, it has been established that situations of clinical ambiguity generate worry for NMPs (Rowbotham et al., 2012). Qualitative studies tend to validate the hypothesis that NMPs are less affected by non-clinical factors, such as patient expectations, to prescribe antibiotics than medical prescribers (Courtenay et al, 2017), and instead rely on protocols and guidelines while making prescribing decisions.

4. Discussion

Prescribing practices are influenced by several factors, and interventions are not likely to address all of them. The findings of this study do not demonstrate a clearly defined pattern of NMP behaviour that influences prescribing behaviour in primary and urgent care facilities, although there are some recurring themes that guide NMP decision-making in these clinical settings. However, this research revealed that non-medical prescribers' decision-making processes were affected by factors such as their belief in their own competence, confidence, the influence of external barriers, and the increased autonomy in their role, as well as the perceived job satisfaction it offered practitioners, during consultations in primary and urgent treatment care facilities. The expanding role of the non-medical prescriber undoubtedly constitutes one of the most substantial potential initiatives aimed at ensuring the sustainability of the NHS at a time of significant change and a drive for improved efficiency (Winter, 2019). In order to maximise the expertise of the workforce in general practise, the acute and community sectors, commissioning authorities will likely be urged to use non-medical prescribers from all professional domains (Weglicki, 2015). NMPs work in primary and urgent care facilities in a variety of locations and capacities. Despite the fact that these prescribers often manage patients and are in charge of 8% of all primary care antibiotic prescriptions, very little study has examined their prescribing practices (Courtenay, 2018). Ambiguity regarding the "proper" prescription behaviour was accentuated by time constraints, limited follow-up capacity, inadequate discussion about prescribing between the physiotherapists and operating physicians, and the anticipation of a therapeutic outcome (Abuzour et al, 2017).

According to the results of this research, participants focused much of their early training on achieving the competency frameworks set by their regulating organisations (Diggle, 2018). As soon as they were given the right to prescribe, however, some NMPs found themselves at a loss for where to go for continuing education in order to keep up with the field or develop their skills (Herklotz et al., 2015) yet little is known about their experiences of prescribing. Drug events and inappropriate prescribing are known to account for around 16% of hospital admissions. With a remit of reducing unplanned admissions to hospital, it is important to understand community matrons' experiences as prescribers and to examine the role of prescribing in fulfilling this agenda with consideration given to the adequacy of the prescribing support available. In this study, a qualitative research design was used and a purposive sample of seven community matrons from the south of England took part in the study. Semi-structured interviews were carried out and analysed using Braun and Clarke's six phases of thematic analysis. The findings show that community matrons consider prescribing-related knowledge to be essential in fulfilling their role and that the ability to prescribe speeds patient access to medicines. In some instances it may be instrumental in preventing hospital admission. Community matrons prescribe a limited range of medicines regularly, while referring to GPs for other prescribing that they consider outside their competency. Community matron prescribers mostly access their support from GPs and this is considered adequate in supporting them in their prescribing role.", "container-title": "British Journal of Community Nursing", "DOI": "10.12968/bjcn.2015.20.5.217", "ISSN": "1462-4753", "issue": "5", "journalAbbreviation": "Br J Community Nurs", "language": "eng", "note": "PMID: 25993369", "page": "217-218, 220-223", "source": "PubMed", "title": "Community matrons' experience as independent prescribers", "volume": "20", "author": [{"family": "Herklotz", "given": "Annie"}, {"family": "Baileff", "given": "Anne"}, {"family": "Latter", "given": "Sue"}], "issued": {"date-parts": [[2015, 5]]}], "schema": "https://github.com/citation-style-language/schema/raw/master/csl-citation.json"} . It was difficult to locate training courses that catered specifically to NMPs, probably due to the tiny size of the NMP population that would benefit from such offerings. This absence of training opportunities was felt most keenly by participants who were unable to discuss prescribing concerns and errors with colleagues and engage in reflective debriefings (Weglicki, 2015). The subsequent lack of self-assurance and consequent reluctance to prescribe was a direct result of the above. In circumstances where they had previously prescribed, they resorted to measures such as referring patients or seeking further guidance (Jarmain, 2022). Lack of communication between professionals can cause inter-professional conflict and effort duplication, which will slow down workflow

and lower the standard of patient care (Graham-Clarke et al., 2019) through extending healthcare professionals' roles. More recent government health service policy focuses on the increased demand and the need for efficiency. This systematic policy review aimed to describe any changes in government policy position and the role that non-medical prescribing plays in healthcare provision.

METHOD: The systematic policy review included policy and consultation documents that describe independent non-medical prescribing. A pre-defined protocol was registered with PROSPERO (CRD42015019786). However, Hindi et al. (2019) highlights that Independent primary care practitioners enjoyed the effective collaboration between NMPs and other members of the healthcare team, which was a significant facilitator to independent prescribing. A team approach to prescribing with peer support and encouragement from medical colleagues helped enhance members' confidence (Cope et al., 2019) Henceforth, participants emphasised how important it is to communicate effectively with the healthcare team, as this facilitated the practitioners' transition into their responsibilities, integration with the larger unit, and ability to function as a cohesive unit (Courtenay et al, 2019).

There was also evidence of a heuristic approach from NMPs, who were shown to depend more on their own intuition and past experiences than on evidence-based pharmacological expertise when addressing clinical problems (Whelehan et al., 2020). According to Abuzour et al. (2017), existing NMPs have indicated a need for continued professional development in the field of pharmacology and medication interactions. This research also discovered that NMPs did not always want to take full responsibility for giving prescriptions for "high-risk" patients at either end of the age range when they didn't know enough and went against protocols, guidelines, formularies, and approved management plans in primary care consultations (Weiss et al., 2015). NMPs also didn't feel comfortable writing prescriptions for people who were on poly-pharmacy prescriptions and suffered from complex co-morbidities. The risk of prescribing without sufficient patient information resulted in a reluctance to prescribe for these category of patients (Armstrong et al., 2021) written in English language. Exclusion criteria: research conducted in secondary care or outside the UK.

Findings:

285 studies were identified; 15 were eligible for critical appraisal. Key themes were: NMP's positive perceptions were autonomy, job satisfaction and colleague support; negative perceptions included risk, lack of continuous professional development (CPD).

4.1. Recommendation

This review revealed a paucity of evidence on the prescribing decisions made by NMPs in primary care and urgent care facilities. The major objective of the examined research was

to examine and explain the prescribing decisions of NMPs. Small sample sizes, primary care settings in the United Kingdom, and the inclusion of virtually entirely nurse prescribers in particular studies all restrict the transferability of the findings of this analysis. This review has conceptualised, using a behavioural perspective, a wide variety of variables and obstacles that require addressing, which may serve as viable targets for behaviour modification strategies to assist non-medical prescribing practice, especially in the primary and urgent care sector (Winter, 2019). Therefore, it is imperative that interventions to improve prescribing practice must be innovated and implemented in a methodical manner, based on grounded data, and guided by evidence-based concepts (Chater, 2020). The most effective method for increasing prescribing confidence involved a blended learning strategy. Ineffective communication between or within primary or secondary care settings is indeed an external obstacle that could also undermine the confidence of non-medical prescribers (Abuzour et al., 2017). Echoing previous research, this study suggests that one way to improve NMPs' competence may be to improve clinical support with opportunities to work alongside established and experienced prescribing practitioners including medical colleagues.

The opportunities and challenges that are faced by NMPs who are employed in primary care are inextricably intertwined, and they are all connected to the concept of continued support for training and continuing professional development (Magowan, 2020). The individual NMP, their employer, educational institutions, and providers all have a responsibility to come up with innovative solutions to address the acknowledged need to update and improve clinical and prescribing procedures that are based on evidence-based processes (Darvishpour et al, 2014). However, the utilisation of evidence-based practise is influenced by a number of different factors that are not immediately correlated to the approach itself. Evidence of poor quality, insufficient evidence, or evidence that contradicts itself all have an impact on the quality of the evidence-base. Hence, the use of evidence in practise has to be redefined in order to make it more malleable and flexible in the future to enhance prescribing behaviour in NMPs (Hubbard, 2019).

5. Conclusion

Although NMP practice is becoming an integral part of healthcare delivery in the United Kingdom and around the world, there is still a substantial dearth of appropriate data on the enablers and barriers experienced by non-medical prescribers, especially in urgent and primary care settings (Beech, 2015). The continuity of care that NMPs provide, particularly in primary care, which frequently has a large turnover of physicians, has had a good influence on the quality

of treatment; clinically incorrect prescriptions are avoided or remedied by specialised non-medical clinicians who have embraced the prescribing role in their field of practise (Diggle, 2018). To ensure that NMP is fully empowered, all facets must be thoroughly scoped before introducing alterations to the scope and practice of non-medical prescribing (Brett and Palmer, 2022). Additionally, there must be clear routes of assistance and supervision for the prescriber, with the objective of fostering support. Clinical supervision and evidence-based practice are effective methods for achieving this objective. Failure to do so may restrict the effective utilization of prescribing skills and contribute to a demotivated workforce (Edwards et al, 2022). In conclusion, this study highlights findings related to NMPs working in primary and urgent care settings, arguing for increased supportive interventions to aid NMPs in developing and upgrading their practice to promote high standards of patient care in these settings.

5.1. Limitations of the Study

There is a plethora of data discussing the influences on NMP prescribing behaviour. However, the authors have chosen only a few qualitative studies to focus on the specific research question, which may have prevented from delving deeper into the topic. In addition, unindexed papers may have been missed from the review inadvertently despite the use of a robust search method to identify relevant publications in the databases using well-considered keyword combinations.

Most of the research focused on prescribing decisions for acute diseases; hence, the findings were not always applicable to individuals with chronic disorders or complex co-morbidities. The studies on prescribing in primary and urgent care settings demonstrates, however, that prescribing decisions are complicated and impacted by several factors, some of which may be contradictory and confound subsequent decision-making. The lack of a comprehensive description of the decision-making process leading to the formulation of a prescription strategy, evaluation, and solution definition is a limitation of the research studies included in this review.

6. Data Availability Statement

All data underlying the results are available as part of the article and no additional source data are required

7. Competing Interests

The authors have no conflict of interest to report

8. Funding

No funding was used to support this research and the preparation of the manuscript

9. Ethics Declaration

The authors have taken sufficient steps to make sure that the work is ethically compliant and adhering to the legal requirements

10. Use of Artificial Intelligence and Adherence to Plagiarism Policy

Part of this work was undertaken as part of a previous UK university continuous professional development program by the first author. If any high Turnitin similarity issues are noticed, the authors declare that it is their own work

References

- Abuzour, A.S., Lewis, P.J. and Tully, M.P. (2018) 'A qualitative study exploring how pharmacist and nurse independent prescribers make clinical decisions', *Journal of Advanced Nursing*, 74(1), pp. 65–74. Available at: <https://doi.org/10.1111/jan.13375>.
- Armstrong, A., Manfrin, A. and Gibson, J. (2021) 'non-medical prescribing in primary care in the United Kingdom: an overview of the current literature', *Journal of Prescribing Practice*, 3(9), pp. 352–361. Available at: <https://doi.org/10.12968/jppr.2021.3.9.352>.
- Atkins, L. et al. (2017) 'A guide to using the Theoretical Domains Framework of behaviour change to investigate implementation problems', *Implementation Science*, 12(1), p. 77. Available at: <https://doi.org/10.1186/s13012-017-0605-9>.
- Aveyard, H. (2018) Doing a Literature Review in Health and Social Care: A Practical Guide. McGraw-Hill Education (UK).
- Aveyard, H. and Bradbury-Jones, C. (2019) 'An analysis of current practices in undertaking literature reviews in nursing: findings from a focused mapping review and synthesis', *BMC medical research methodology*, 19(1), p. 105. Available at: <https://doi.org/10.1186/s12874-019-0751-7>.
- Aveyard, H., Payne, S. and Preston, N. (2021) A Postgraduate's Guide to Doing a Literature Review in Health and Social Care, 2e. McGraw-Hill Education (UK).
- Beauchamp, T.L. and Childress, J.F. (2019) Principles of Biomedical Ethics. Oxford University Press.
- Beech, E. (2015) 'Promoting appropriate antimicrobial prescribing in primary care'.
- Berry, D., Bradlow, A. and Courtenay, M. (2008) 'Patients' attitudes towards, and information needs in relation

- to, nurse prescribing in rheumatology', *Journal of Clinical Nursing*, 17(2), pp. 266–273. Available at: <https://doi.org/10.1111/j.1365-2702.2007.01980.x>.
- Bettany-Saltikov, J. and McSherry, R. (2016) How to do a Systematic Literature Review in Nursing: A step-by-step guide (2nd Edition).
- Booth, A. et al. (2015) What is the evidence for the effectiveness, appropriateness and feasibility of group clinics for patients with chronic conditions? A systematic review. Southampton (UK): NIHR Journals Library (*Health Services and Delivery Research*). Available at: <http://www.ncbi.nlm.nih.gov/books/NBK333454/>.
- Booth, A. (2016) 'Searching for qualitative research for inclusion in systematic reviews: a structured methodological review', *Systematic Reviews*, 5, p. 74. Available at: <https://doi.org/10.1186/s13643-016-0249-x>.
- Booth, A. and Carroll, C. (2015) 'Systematic searching for theory to inform systematic re-views: is it feasible? Is it desirable?', *Health Information and Libraries Journal*, 32(3), pp. 220–235. Available at: <https://doi.org/10.1111/hir.12108>.
- Braun, V. and Clarke, V. (2013) Successful Qualitative Research: A Practical Guide for Beginners. SAGE Publications.
- Brett, E.A. and Palmer, M. (2022) 'The influence of non-medical prescribers on antimicrobial stewardship: a national evaluation of the prescribing of antibiotics by non-medical prescribers in England from 2016 to 2021 (part 1)', *Journal of Prescribing Practice*, 4(10), pp. 452–462. Available at: <https://doi.org/10.12968/jppr.2022.4.10.452>.
- Brún, C.D. (2013) 'A key step in the information production process'.
- Bryant-Lukosius, D. et al. (2016) 'Framework for Evaluating the Impact of Advanced Practice Nursing Roles', *Journal of Nursing Scholarship: An Official Publication of Sigma Theta Tau International Honor Society of Nursing*, 48(2), pp. 201–209. Available at: <https://doi.org/10.1111/jnu.12199>.
- Buccheri, R.K. and Sharifi, C. (2017) 'Critical Appraisal Tools and Reporting Guidelines for Evidence-Based Practice', *Worldviews on Evidence-Based Nursing*, 14(6), pp. 463–472. Available at: <https://doi.org/10.1111/wvn.12258>.
- Chater, N., 2020. Facing up to the uncertainties of COVID-19. *Nature Human Behaviour*, [online] 4(5), pp.439–439. Available at: <https://doi.org/10.1038/s41562-020-0865-2>.
- Claydon, L.S. (2015) 'Rigour in quantitative research', *Nursing Standard (Royal College of Nursing (Great Britain): 1987)*, 29(47), pp. 43–48. Available at: <https://doi.org/10.7748/ns.29.47.43.e8820>.
- Conner, M. and Norman, P. (2015) Predicting and Changing Health Behaviour: Research and Practice with Social Cognition Models. Open University Press.
- Cope, L.C., Tully, M.P. and Hall, J. (2020) 'An exploration of the perceptions of non-medical prescribers, regarding their self-efficacy when prescribing, and their willingness to take responsibility for prescribing decisions', *Research in Social and Administrative Pharmacy*, 16(2), pp. 249–256. Available at: <https://doi.org/10.1016/j.sapharm.2019.05.013>.
- Courtenay, M. et al. (2017) 'Overview of the uptake and implementation of non-medical prescribing in Wales: a national survey', *BMJ open*, 7(9), p. e015313. Available at: <https://doi.org/10.1136/bmjopen-2016-015313>.
- Courtenay, M. (2018) 'An overview of developments in nurse prescribing in the UK', *Nursing Standard (Royal College of Nursing (Great Britain): 1987)*, 33(1), pp. 40–44. Available at: <https://doi.org/10.7748/ns.2018.e11078>.
- Courtenay, M. et al. (2019) 'Examining influences on antibiotic prescribing by nurse and pharmacist prescribers: a qualitative study using the Theoretical Domains Framework and COM-B', *BMJ Open*, 9(6), p. e029177. Available at: <https://doi.org/10.1136/bmjopen-2019-029177>.
- Courtenay, M. and Griffiths, M. (2022) Independent and Supplementary Prescribing: An Essential Guide. Cambridge University Press.
- Cousins, R. and Donnell, C. (2012) 'Nurse prescribing in general practice: a qualitative study of job satisfaction and work-related stress', *Family Practice*, 29(2), pp. 223–227. Available at: <https://doi.org/10.1093/fampra/cmr077>.
- Creswell, J.W. et al. (2007) 'Qualitative Research Designs: Selection and Implementation', *The Counseling Psychologist*, 35(2), pp. 236–264. Available at: <https://doi.org/10.1177/001100006287390>.
- Darvishpour, A., Joolaee, S. and Cheraghi, M.A. (2014) 'A meta-synthesis study of literature review and systematic review published in nurse prescribing', *Medical Journal of the Islamic Republic of Iran*, 28, p. 77. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4219909/> (Accessed: 8 January 2023).
- Day, M. (2005) 'UK doctors protest at extension to nurses' prescribing powers', *BMJ: British Medical Journal*, 331(7526), p. 1159. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4219909/>

- nlm.nih.gov/pmc/articles/PMC1285084/ (Accessed: 6 January 2023).
- Department of Health (2006) 'Improving patients' access to medicines: a guide to implementing nurse and pharmacist independent prescribing within the NHS in England'. Available at: http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_4133743.
- Diggle, J. (2018) 'The Diabetes Multidisciplinary Team: How do nurse prescribers' demonstrate prescribing proficiency? - DiabetesontheNet', 31 May. Available at: <https://diabetesonthenet.com/diabetes-primary-care/nurse-prescribers-prescribing-proficiency/>.
- Edwards, J., Coward, M. and Carey, N. (2022) 'Barriers and facilitators to implementation of non-medical independent prescribing in primary care in the UK: a qualitative systematic review', *BMJ open*, 12(6), p. e052227. Available at: <https://doi.org/10.1136/bmjopen-2021-052227>.
- Franklin, P.M. (2017) Non-medical Prescribing in the United Kingdom. Springer.
- Gerrish, K. and Lacey, A. (2013) The Research Process in Nursing. John Wiley & Sons.
- Graham-Clarke, E. et al. (2019) 'Non-medical prescribing in the United Kingdom National Health Service: A systematic policy review', *PLoS One*, 14(7), p. e0214630. Available at: <https://doi.org/10.1371/journal.pone.0214630>.
- Graham-Clarke, E., Rushton, A. and Marriott, J. (2021) 'A Delphi study to explore and gain consensus regarding the most important barriers and facilitators affecting physiotherapist and pharmacist non-medical prescribing', *PLOS ONE*, 16(2), p. e0246273. Available at: <https://doi.org/10.1371/journal.pone.0246273>.
- Grant, M.J. and Booth, A. (2009) 'A typology of reviews: an analysis of 14 review types and associated methodologies', *Health Information and Libraries Journal*, 26(2), pp. 91–108. Available at: <https://doi.org/10.1111/j.1471-1842.2009.00848.x>.
- Greenhalgh, T. (2010) How to Read a Paper: The Basics of Evidence-Based Medicine. John Wiley & Sons.
- Griffith, R. and Tengnah, C. (2014) Law and Professional Issues in Nursing. Learning Mat-ters.
- Hacking, S. and Taylor, J. (2010) 'An evaluation of the scope and practice of Non-Medical Prescribing', School of Nursing & Caring Sciences University of Central Lancashire [Pre-print].
- Harden, A. and Gough, D. (2012) 'Quality and relevance appraisal', in. Sage Publications.
- Harvey, G. and Kitson, A. (2015) Implementing Evidence-Based Practice in Healthcare: A Facilitation Guide. Routledge.
- Health Education England (2015) The future of primary care Creating teams for tomorrow. Primary Care Workforce Commission. Available at: <https://www.oxfordhealth.nhs.uk/library/wp-content/uploads/sites/3/2018/03/The-Future-of-Primary-Care-report-HEE.pdf>.
- Herklotz, A., Baileff, A. and Latter, S. (2015) 'Community matrons' experience as independent prescribers', *British Journal of Community Nursing*, 20(5), pp. 217–218, 220–223. Available at: <https://doi.org/10.12968/bjcn.2015.20.5.217>.
- Higgins, J. and Thomas, J. (2022) Cochrane Handbook for Systematic Reviews of Interventions. Available at: <https://training.cochrane.org/handbook/current>.
- Hindi, A.M.K. et al. (2019) 'Independent prescribing in primary care: A survey of patients', prescribers' and colleagues' perceptions and experiences', *Health & Social Care in the Community*, 27(4), pp. e459–e470. Available at: <https://doi.org/10.1111/hsc.12746>.
- Holly, C., Salmond, S.W. and Saimbert, M. (2011) Comprehensive Systematic Review for Advanced Nursing Practice. Springer Publishing Company.
- Hubbard, K. (2019) 'My experience as a non-medical prescriber from 2012–2019: are we still facing the same problems?', *Journal of Prescribing Practice*, 1(6), pp. 302–305. Available at: <https://doi.org/10.12968/jppr.2019.1.6.302>.
- Jarmain, S. (2022) 'Describing prescribing identities: a qualitative study exploring non-medical prescriber identity', *Journal of Prescribing Practice*, 4(7), pp. 300–307. Available at: <https://doi.org/10.12968/jppr.2022.4.7.300>.
- Kumar, R. (2010) Research Methodology: A Step-by-Step Guide for Beginners. SAGE.
- Lennon, R. and Fallon, A. (2018) 'The experiences of being a registered nurse prescriber within an acute service setting', *Journal of Clinical Nursing*, 27(3–4), pp. e523–e534. Available at: <https://doi.org/10.1111/jocn.14087>.
- Lim, A.G., North, N. and Shaw, J. (2017) 'Navigating professional and prescribing boundaries: Implementing nurse prescribing in New Zealand', *Nurse Education in Practice*, 27, pp. 1–6. Available at: <https://doi.org/10.1016/j.nepr.2017.08.009>.
- Maddox, C. et al. (2016) 'Factors influencing nurse and pharmacist willingness to take or not take responsibility for non-medical prescribing', *Research*

- in social & administrative pharmacy: RSAP*, 12(1), pp. 41–55. Available at: <https://doi.org/10.1016/j.sapharm.2015.04.001>.
- Magowan, J. (2020) ‘Barriers and enablers to nurse prescribing in primary care’, *Journal of Prescribing Practice*, 2(3), pp. 142–146. Available at: <https://doi.org/10.12968/jppr.2020.2.3.142>.
- Mays, N., Pope, C. and Popay, J. (2005) ‘Systematically reviewing qualitative and quantitative evidence to inform management and policy-making in the health field’, *Journal of Health Services Research & Policy*, 10 Suppl 1, pp. 6–20. Available at: <https://doi.org/10.1258/1355819054308576>.
- McHugh, Á. et al. (2020) ‘non-medical prescribers: prescribing within practice’, *Journal of Prescribing Practice*, 2(2), pp. 68–77. Available at: <https://doi.org/10.12968/jppr.2020.2.2.68>.
- McIntosh, T. et al. (2016) ‘Influences on prescribing decision-making among non-medical prescribers in the United Kingdom: systematic review’, *Family Practice*, 33(6), pp. 572–579. Available at: <https://doi.org/10.1093/fampra/cmw085>.
- Moher, D. et al. (2009) ‘Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement’, *PLoS medicine*, 6(7), p. e1000097. Available at: <https://doi.org/10.1371/journal.pmed.1000097>.
- Moule, P., Aveyard, H. and Goodman, M. (2016) *Nursing Research: An Introduction*. SAGE.
- Munn, Z., Tufanaru, C. and Aromataris, E. (2014) ‘JBI’s Systematic Reviews: Data Extraction and Synthesis’, *AJN The American Journal of Nursing*, 114(7), p. 49. Available at: <https://doi.org/10.1097/01.NAJ.0000451683.66447.89>.
- Nelson, P.A. et al. (2019) ‘Skill-mix change in general practice: a qualitative comparison of three “new” non-medical roles in English primary care’, *British Journal of General Practice*, 69(684), pp. e489–e498. Available at: <https://doi.org/10.3399/bjgp19X704117>.
- NHS England (2013) NHS England » Updated NHS Constitution published. Available at: <https://www.england.nhs.uk/2013/03/nhs-constitution/>.
- NHS England (2014) ‘Five Year Forward View’. Available at: <https://www.england.nhs.uk/wp-content/uploads/2014/10/5yfv-web.pdf>.
- NHS England (2018) NHS England » Non-medical prescribing by allied health professionals. Available at: <https://www.england.nhs.uk/ahp/med-project/>.
- NHS England (2019) ‘The NHS Long Term Plan’. f. Available at: <https://www.longtermplan.nhs.uk/>
- <wp-content/uploads/2019/08/nhs-long-term-plan-version-1.2.pdf>.
- NICE (2012) National Prescribing Centre/National Institute for Health and Clinical Excellence (NICE). The Prescribing Competency Framework for all prescribers. Available at: <https://www.rpharms.com/resources/frameworks/prescribers-competency>.
- Nuttall, D. (2018) ‘Nurse prescribing in primary care: a Meta synthesis of the literature’, *Primary Health Care Research & Development*, 19(1), pp. 7–22. Available at: <https://doi.org/10.1017/S1463423617000500>.
- Offredy, M. and Vickers, P. (2010) *Developing a Healthcare Research Proposal: An Inter-active Student Guide*. John Wiley & Sons.
- Parahoo, K. (2014) *Nursing Research: Principles, Process and Issues*. Bloomsbury Publishing.
- Paré, G. et al. (2015) ‘Synthesizing information systems knowledge: A typology of literature reviews’, *Information & Management*, 52(2), pp. 183–199. Available at: <https://doi.org/10.1016/j.im.2014.08.008>.
- Park, J. et al. (2018) ‘To What Extent Are State Scope of Practice Laws Related to Nurse Practitioners’ Day-to-Day Practice Autonomy?’, *Medical care research and review: MCRR*, 75(1), pp. 66–87. Available at: <https://doi.org/10.1177/1077558716677826>.
- Patton, M.Q. (1999) ‘Enhancing the quality and credibility of qualitative analysis’, *Health Services Research*, 34(5 Pt 2), pp. 1189–1208.
- Pearce, L. (2016) The long road to full prescribing powers. RCNi. Available at: <https://rcni.com/nursing-standard/features/long-road-full-prescribing-powers-56726> (Accessed: 6 January 2023).
- Polit, D.F. and Beck, C.T. (2017) *Nursing Research: Generating and Assessing Evidence for Nursing Practice*. Wolters Kluwer Health.
- Popay, J. et al. (2006) ‘Guidance on the Conduct of Narrative Synthesis in Systematic Re-views’. Available at: <https://www.lancaster.ac.uk/media/lancaster-university/content-assets/documents/fhm/dhr/chir/NSsynthesisguidanceVersion1-April2006.pdf>.
- Purssell, E. (2020) ‘Can the Critical Appraisal Skills Programme checklists be used along-side Grading of Recommendations Assessment, Development and Evaluation to improve transparency and decision-making?’, *Journal of Advanced Nursing*, 76(4), pp. 1082–1089. Available at: <https://doi.org/10.1111/jan.14303>.
- Pussegoda, K., Turner, L., Garrity, C., Mayhew, A., Skidmore, B., Stevens, A., Boutron, I., Sarkis-Onofre, R., Bjerre, L.M., Hróbjartsson, A., Altman, D.G. and Moher, D., 2017. Identifying approaches for assessing

- methodological and reporting quality of systematic reviews: a descriptive study. *Systematic Reviews*, [online] 6(1). <https://doi.org/10.1186/s13643-017-0507-6>.
- Reeve, J. et al. (2018) 'Identifying enablers and barriers to individually tailored prescribing: a survey of healthcare professionals in the UK', *BMC Family Practice*, 19(1), p. 17. Available at: <https://doi.org/10.1186/s12875-017-0705-2>.
- Ritchie, J. et al. (2013) Qualitative Research Practice: A Guide for Social Science Students and Researchers. SAGE.
- Robertson, D. (2022) 'Impact of non-medical prescribing', *Journal of Prescribing Practice*, 4(1), pp. 12–13. Available at: <https://doi.org/10.12968/jppr.2022.4.1.12>.
- Rowbotham, S. et al. (2012) 'Challenges to nurse prescribers of a no-antibiotic prescribing strategy for managing self-limiting respiratory tract infections', *Journal of Advanced Nursing*, 68(12), pp. 2622–2632. Available at: <https://doi.org/10.1111/j.1365-2648.2012.05960.x>.
- Royal Pharmaceutical Society (2016) Prescribing Competency Framework | RPS. Available at: <https://www.rpharms.com/resources/frameworks/prescribers-competency-framework>.
- Smith, A., Latter, S. and Blenkinsopp, A. (2014) 'Safety and quality of nurse independent prescribing: a national study of experiences of education, continuing professional development clinical governance', *Journal of Advanced Nursing*, 70(11), pp. 2506–2517. Available at: <https://doi.org/10.1111/jan.12392>.
- Smith, J. and Noble, H. (2016) 'Reviewing the literature', *Evidence-Based Nursing*, 19(1), pp. 2–3. Available at: <https://doi.org/10.1136/eb-2015-102252>.
- Stewart, D. et al. (2017) 'Future perspectives on nonmedical prescribing', *Therapeutic Advances in Drug Safety*, 8(6), pp. 183–197. Available at: <https://doi.org/10.1177/2042098617693546>.
- Stewart, D., MacLure, K. and George, J. (2012) 'Educating nonmedical prescribers', *British Journal of Clinical Pharmacology*, 74(4), pp. 662–667. Available at: <https://doi.org/10.1111/j.1365-2125.2012.04204.x>.
- Sun, N. et al. (2020) 'A qualitative study on the psychological experience of caregivers of COVID-19 patients', *American Journal of Infection Control*, 48(6), pp. 592–598. Available at: <https://doi.org/10.1016/j.ajic.2020.03.018>.
- The King's Fund (2019) The NHS long-term plan explained, The King's Fund. Available at: <https://www.kingsfund.org.uk/publications/nhs-long-term-plan-explained>.
- Thomas, D. and Hodges, I.D. (2010) Designing and Managing Your Research Project: Core Skills for Social and Health Research. SAGE.
- Weeks, G. et al. (2016) 'Non-medical prescribing versus medical prescribing for acute and chronic disease management in primary and secondary care', *The Cochrane Database of Systematic Reviews*, 11(11), p. CD011227. Available at: <https://doi.org/10.1002/14651858.CD011227.pub2>.
- Weglicki, R.S., Reynolds, J. and Rivers, P.H. (2015) 'Continuing professional development needs of nursing and allied health professionals with responsibility for prescribing', *Nurse Education Today*, 35(1), pp. 227–231. Available at: <https://doi.org/10.1016/j.nedt.2014.08.009>.
- Weiss, M.C. et al. (2015) 'Medication decision making and patient outcomes in GP, nurse and pharmacist prescriber consultations', *Primary Health Care Research & Development*, 16(5), pp. 513–527. Available at: <https://doi.org/10.1017/S146342361400053X>.
- Whelahan, D.F., Conlon, K.C. and Ridgway, P.F. (2020) 'Medicine and heuristics: cognitive biases and medical decision-making', *Irish Journal of Medical Science*, 189(4), pp. 1477–1484. Available at: <https://doi.org/10.1007/s11845-020-02235-1>.
- WHO (2021) Antibiotic resistance. Available at: <https://www.who.int/news-room/fact-sheets/detail/antibiotic-resistance>.
- Williamson, G.R. and Whittaker, A. (2017) Succeeding in Literature Reviews and Research Project Plans for Nursing Students. Learning Matters.
- Williamson, V., Murphy, D. and Greenberg, N. (2020) 'COVID-19 and experiences of moral injury in front-line key workers', *Occupational Medicine (Oxford, England)*, 70(5), pp. 317–319. Available at: <https://doi.org/10.1093/occmed/kqaa052>.
- Wilson, N. et al. (2020) 'Limited pharmaceuticalisation: a qualitative case study of physiotherapist prescribing practices in an NHS Trust in England following the expansion of non-medical prescribing in the UK', *Sociology of Health & Illness*, 42(3), pp. 643–659. Available at: <https://doi.org/10.1111/1467-9566.13050>.
- Winter, G. (2019) 'The NHS Long Term Plan', *Journal of Prescribing Practice*, 1(3), pp. 114–114. Available at: <https://doi.org/10.12968/jppr.2019.1.3.114>.



Journal of Multidisciplinary Research in Healthcare

Chitkara University, Saraswati Kendra, SCO 160-161, Sector 9-C,
Chandigarh, 160009, India

Volume 8, Issue 2

April 2022

ISSN 2393-8536

Copyright: [© 2022 Mithun Thampi and Biju Mathew] This is an Open Access article published in Journal of Multidisciplinary Research in Healthcare (J. Multidiscip. Res. Healthcare) by Chitkara University Publications. It is published with a Creative Commons Attribution- CC-BY 4.0 International License. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.
